

**STABILITY OF FUNCTIONAL INEQUALITIES WITH 3K-VARIABLE BASED
ON JORDAN-VON NEUMANN TYPE ADDITIVE FUNCTIONAL EQUATIONS
IN BANACH SPACE****Ly Van An*****Faculty of Mathematics Teacher Education, Tay Ninh University, Tay Ninh, Vietnam.****Corresponding Author:***lyvanan145@gmail.com, lyvananvietnam@gmail.com.***Abstract**

In this paper, we study to solve the Cauchy, Jensen and Cauchy-Jensen additive function inequalities with 3k-variables related to Jordan-von Neumann type in Banach space. These are the main results of this paper.

Keywords

Normed Spaces; Banach Space; Generalized Hyers-Ulam-Rassias Stability Jordan-von Neumann-Type Additive Functional Equation; Cauchy, Jensen Additive Function Inequalities.

1. INTRODUCTION

Let X and Y be a normed spaces on the same field K , and $f : X \rightarrow Y$ be a mapping. We use the notation $\|\cdot\|_X$ and $\|\cdot\|_Y$ for corresponding the norms on X and Y . In this paper, we investigate additive functional inequalities associated with Jordan-Von Neumann type additive functional equation when X is a normed space with norm $\|\cdot\|_X$ and that Y is a Banach space with norm $\|\cdot\|_Y$. In fact, when X is a normed space with norm $\|\cdot\|_X$ and that Y is a Banach space with norm $\|\cdot\|_Y$ we solve and prove the Hyers – Ulam – Rassias type stability of following additive functional inequalities

$$\left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_Y \leq \left\| 2kf \left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k} \right) \right\|_Y, \quad (1.1)$$

and

$$\left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_Y \leq \left\| f \left(\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j \right) \right\|_Y, \quad (1.2)$$

final

$$\left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + 2k \sum_{j=1}^k f(z_j) \right\|_Y \leq \left\| 2kf \left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j}{2k} + \sum_{j=1}^k z_j \right) \right\|_Y. \quad (1.3)$$

The study the stability of generalized additive functional inequalities associated with jordan-von neumann type additive functional equation originated from a question of S.M. Ulam[1], concerning the stability of group homomorphisms.

Let $(G, *)$ be a group and let (G_0, \circ, d) be a metric group with metric $(d(\cdot, \cdot))$. Given $\epsilon > 0$, does there exist a $\delta > 0$ such that if $f : G \rightarrow G_0$ satisfies

$$d(f(x * y), f(x) \circ f(y)) < \delta, \forall x \in G$$

then there is a homomorphism $h : G \rightarrow G_0$

$$d(f(x), h(x)) < \epsilon, \forall x \in G$$

The concept of stability for a functional equation arises when we replace functional equation by an inequality which acts as a perturbation of the equation. Thus the stability question of functional equations is that how do the solutions of the inequality differ from those of the given function equation? Hyers gave a first affirmative answer to the question of Ulam as follows: In 1941 D. H. Hyers [2] Let $\epsilon \geq 0$ and let $f : E_1 \rightarrow E_2$ be a mapping between Banach space

$$\|f(x + y) - f(x) - f(y)\| \leq \epsilon, \quad (1.4)$$

for all $x, y \in E_1$ and some $\epsilon \geq 0$. It was shown that the limit

$$T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad (1.5)$$

exists for all $x \in E_1$ and that $T : E_1 \rightarrow E_2$ is that unique additive mapping satisfying

$$\|f(x) - T(x)\| \leq \epsilon, \forall x \in E_1. \quad (1.6)$$

Next in 1978 Th. M. Rassias [3] provided a generalization of Hyers' Theorem which allows the Cauchy difference to be unbounded:

Consider E, E_0 to be two Banach spaces, and let $f : E \rightarrow E_0$ be a mapping such that $f(tx)$ is continuous in t for each fixed x . Assume that there exist $\theta \geq 0$ and $p \in [0, 1)$, $\epsilon > 0$ Such that

$$\|f(x + y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p), \forall x, y \in E. \quad (1.7)$$

Where ϵ and p is constants with $\epsilon > 0$ and $p < 1$. Then the limit

$$T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad (1.8)$$

there exists a unique linear $L : \mathbf{E} \rightarrow \mathbf{E}'$ satisfies

$$\|f(x) - L(x)\| \leq \frac{2\epsilon}{2-2^p} \|x\|^p, x \in \mathbf{E}. \quad (1.9)$$

If $p < 0$, then inequality (1.7) holds for $x, y \neq 0$ and (1.9) for $x \neq 0$

We notice that in Rassias' functional inequality (1.7) Mathematicians around the world such as [4],[5] as well as Rassias have asserted that the inequality (1.7) no longer holds true when $p = 1$ from the assertion that gave rise to the idea to generalize the generalized functional equation Hyers- Ulam more specifically.

Thus, to replace the non-existent condition mentioned above, Mathematician Rassias

[2] has given the following specific conditions: $\|x\|^p + \|y\|^p$ by $\|x\|^p \|y\|^p$ for $p, q \in \mathbb{R}$ with $p+q = 1$.

for all $x, y \in \mathbf{E}$ Gāvruta[6] provided a further generalization of Rassias theorem. During the last two decades, a number of papers and research monographs have been published on various generalizations and applications of the generalized Hyers-Ulam stability to a number of functional equations and mappings.

Afterward Gila'ny [7] showed that if f satisfies the functional inequality

$$\left\| 2f(x) + 2f(y) - f(xy^{-1}) \right\| \leq \|f(xy)\| \quad (1.10)$$

f satisfies the Jordan-von Newman functional equation

$$2f(x) + 2f(y) = f(xy) + f(xy^{-1}) \quad (1.11)$$

Then, mathematicians in the world proved to extend the functional inequality (1.11) as [7]-[17].In addition, mathematicians have developed the achievements of their predecessors who have built mathematical models from advanced to modern mathematics, especially functional equations applied on function spaces to Unlocking means connecting with other Maths. [3]-[35]Recently, the authors studied the Hyers-Ulam-Rassias type stability for the following functional inequalities (see [31],[32],[34])

$$\left\| f(x) + f(y) + f(z) \right\| \leq \left\| k \left(f\left(\frac{x+y+z}{k}\right) \right) \right\|, |k| < 3, \quad (1.12)$$

$$\left\| f(x_1) + f(x_2) + \dots + f(x_n) \right\| \leq \left\| kf\left(\frac{x_1+x_2+\dots+x_n}{k}\right) \right\|, |n| > |k|, \quad (1.13)$$

$$\left\| \sum_{j=1}^n f(x_j) + \sum_{j=1}^n f\left(\frac{x_{n+j}}{n}\right) \right\| \leq \left\| kf\left(\frac{\sum_{j=1}^n x_j}{k} + \frac{\sum_{j=1}^n x_{n+j}}{n \cdot k}\right) \right\|, |n| > |k|. \quad (1.14)$$

In banach spaces.

In this paper, we solve and proved the Hyers-Ulam- Rassias type stability for functional inequalities (1.1), (1.2) and (1.3) ie the functional inequalities with $3k$ -variables. Under suitable assumptions on spaces \mathbf{X} and \mathbf{Y} , we will prove that the mappings satisfying the functional inequalities (1.1), (1.2) and (1.3). Thus, the results in this paper are generalization of those in [21],[31],[32],[34] for functional inequalities with $3k$ -variables.

The paper is organized as follows:

In section preliminaries we remind some basic notations in such as Solutions of the inequalities.

Section:3 The basis for building solutions for functional inequalities related to the type of Jordan-Neuman additive functional equations

Section:4 Establishing solutions to functional inequality (1.1) related to the type of Jensen additive functional equation

Section:5 Establishing solutions to functional inequality (1.2) related to the type of Cauchy additive functional equation.

Section:6 Establishing solutions to functional inequality (1.3) related to the type of Cauchy-Jensen additive functional equation.

2. PRELIMINARIES

2.1. **Solutions of the inequalities.** The functional equation

$$f(x+y) = f(x) + f(y)$$

is called the Cauchy equation. In particular, every solution of the Cauchy equation is said to be an *additive mapping*.

The functional equation

$$f\left(\frac{x+y}{2}\right) = \frac{1}{2}f(x) + \frac{1}{2}f(y)$$

is called the Jensen equation. In particular, every solution of the Jensen equation is said to be a Jensen *additive mapping*.

The functional equation

$$2f\left(\frac{x+y}{2} + z\right) = f(x) + f(y) + 2f(z)$$

is called the Cauchy-Jensen equation. In particular, every solution of the Cauchy-Jensen equation is said to be a Jensen-Cauchy *additive mapping*.

3. THE BASIS FOR BUILDING SOLUTIONS FOR FUNCTIONAL INEQUALITIES RELATED TO THE TYPE OF JORDAN-NEUMAN ADDITIVE FUNCTIONAL EQUATIONS

The basis for building solutions for functional inequalities related to the type of Jordan-Neuman additive functional equations. Now, we first study the solutions of (1.1), (1.2) and (1.3). Note that for this inequalities, X is a normed space with norm $\|\cdot\|_X$ and that Y is a Banach space with norm $\|\cdot\|_Y$. Under this setting, we can show that the mappings satisfying (1.1), (1.2) and (1.3) is additive.

Here we assume that G is a 3k-divisible abelian group.

Proposition 3.1. Suppose $f: X \rightarrow Y$ be a mapping such that

$$\left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_Y \leq \left\| 2kf\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k}\right) \right\|_Y \quad (3.1)$$

for all $x_j, y_j, z_j \in X$ for all $j = 1 \rightarrow n$ then f is additive.

Proof. Assume that $f: X \rightarrow Y$ satisfies (3.1).

We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $0, \dots, 0, 0, \dots, 0, 0, \dots, 0$ in (3.1), we have $f(0) = 0$.

Next We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $x, 0, \dots, 0, -x, 0, \dots, 0, 0, \dots, 0$ in (3.1), we have

$$\|f(x) + f(-x)\|_Y \leq \|2nf(0)\|_Y \quad (3.2)$$

, for all $x \in X$.

Hence $f(x) = -f(-x)$, $\forall x \in X$

Next We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $x, 0, \dots, 0, y, 0, \dots, 0, -x - y, \dots, 0$ in (3.1), we have

$$\|f(x) + f(y) - f(x+y)\|_Y = \|f(x) + f(y) + f(-x-y)\|_Y \leq \|2nf(0)\|_Y = 0 \quad (3.3)$$

for all $x, y \in X$. It follows that $f(x+y) = f(x) + f(y)$ This completes the proof.

Proposition 3.2. $f: X \rightarrow Y$ be a mapping such a that

$$\left\| \sum_{j=1}^n f(x_j) + \sum_{j=1}^n f(y_j) + \sum_{j=1}^n f(z_j) \right\|_Y \leq \left\| f\left(\sum_{j=1}^n x_j + \sum_{j=1}^n y_j + \sum_{j=1}^n z_j\right) \right\|_Y \quad (3.4)$$

for all $x_j, y_j, z_j \in X$ for all $j = 1 \rightarrow n$ then f is additive.

Proof. Assume that $f: X \rightarrow Y$ satisfies (3.4).

We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $0, \dots, 0, 0, \dots, 0, 0, \dots, 0$ in (3.4), we have $f(0) = 0$.

Next We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $(x, \dots, 0, -x, \dots, 0, 0, \dots, 0)$ in (3.4), we have

$$\|f(x) + f(-x)\|_Y \leq \|f(0)\|_Y \quad (3.5)$$

for all $x \in X$.

Hence. $f(x) = -f(-x)$, $\forall x \in X$

Next We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $(0, \dots, 0, y, 0, \dots, 0, -x - y, \dots, 0)$ in (3.4), we have

$$\|f(x) + f(y) - f(x+y)\|_Y = \|f(x) + f(y) + f(-x-y)\|_Y \leq \|f(0)\|_Y = 0 \quad (3.6)$$

for all $x, y \in X$. It follows that $f(x+y) = f(x) + f(y)$ This completes the proof.

Proposition 3.3. $f: G \rightarrow Y$ be a mapping such that

$$\left\| \sum_{j=1}^n f(x_j) + \sum_{j=1}^n f(y_j) + 2n \sum_{j=1}^n f(z_j) \right\|_Y \leq \left\| 2nf \left(\frac{\sum_{j=1}^n x_j + \sum_{j=1}^n y_j}{2n} + \sum_{j=1}^n z_j \right) \right\|_Y \quad (3.7)$$

for all $x_j, y_j, z_j \in X$ for all $j = 1 \rightarrow n$ then f is additive.

Proof. Assume that $f: G \rightarrow Y$ satisfies (3.7).

We replacing $(x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n)$ by $(0, \dots, 0, 0, \dots, 0, 0, \dots, 0)$ in (3.7), we have we get

$$\|(2n^2 + 2n)f(0)\| \leq \|2nf(0)\|_Y \quad (3.8)$$

. So $f(0) = 0$

Next We replacing $(x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n)$ by $(x, 0, \dots, 0, -x, 0, \dots, 0, 0, \dots, 0)$ in (3.7), we have

$$\|f(x) + f(-x)\|_Y \leq \|2nf(0)\|_Y \quad (3.9)$$

for all $x \in X$.

Hence. $f(x) = -f(-x)$, $\forall x \in X$

Next We replacing $(x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n)$ by $(-2nz, 0, \dots, 0, 0, \dots, 0, z, 0, \dots, 0)$ in (3.7), we have

$$\|f(-2nz) + 2nf(z)\|_Y \leq \|2nf(0)\|_Y \quad (3.10)$$

for all $x \in X$.

Thus $f(2nz) = 2nf(z)$, $\forall z \in G$.

Next We replacing $(x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n)$ by $(x_1, \dots, x_n, y_1, \dots, y_n, -\frac{x_1+y_1}{2n}, \dots, -\frac{x_n+y_n}{2n})$ in (3.7), we have

$$\begin{aligned} & \left\| \sum_{j=1}^n f(x_j) + \sum_{j=1}^n f(y_j) - \sum_{j=1}^n f(x_j + y_j) \right\|_Y \\ & \quad \left\| \sum_{j=1}^n f(x_j) + \sum_{j=1}^n f(y_j) + 2n \sum_{j=1}^n f\left(-\frac{x_j + y_j}{2n}\right) \right\|_Y \\ & \quad \leq \|2nf(0)\|_Y \end{aligned} \quad (3.11)$$

$\forall x_1, \dots, x_k, y_1, \dots, y_k, -\frac{x_1+y_1}{2n}, \dots, -\frac{x_k+y_k}{2n} \in \mathbf{G}_{\text{Thus}}$

$$\sum_{j=1}^n f(x_j) + \sum_{j=1}^n f(y_j) - \sum_{j=1}^n f(x_j + y_j) = 0 \quad (3.12)$$

Next put $x = x_j, y = y_j$ for all $j = 1 \rightarrow n$ in (3.12), we have

$$f(x + y) = f(x) + f(y)$$

for all $x, y \in \mathbf{G}$. It follows that f is an additive mapping and the proof is complete.

4. Establishing solutions to functional inequality (1.1) related to the type of Jensen additive functional equation

Now, we first study the solutions of (1.1). Note that for this inequality, \mathbf{X} is a normed space with norm and that \mathbf{Y} is a Banach space with norm. Under this setting, we can show that the mappings satisfying (1.1) is Jensen additive. These results are given in the following.

Theorem 4.1. Suppose $q > 1$, θ be non-negative real and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned} & \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} \\ & \leq \left\| 2kf\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k}\right) \right\|_{\mathbf{Y}} \\ & + \theta \left(\sum_{j=1}^k \|x_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|y_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|z_j\|_{\mathbf{X}}^q \right) \end{aligned} \quad (4.1)$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbf{Y}} \leq \frac{(2k)^q + 2k}{(2k)^q - 2k} \theta \|x\|_{\mathbf{X}}^q. \quad (4.2)$$

for all $x \in \mathbf{X}$.

Proof. Assume that $f: \mathbf{X} \rightarrow \mathbf{Y}$ satisfies (4.1).

We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $(x, \dots, x, x, \dots, x, -2kx, 0, \dots, 0)$ in (4.1), we have

$$\left\| 2kf(x) - f(2kx) \right\|_{\mathbf{Y}} = \left\| 2kf(x) + f(-2kx) \right\|_{\mathbf{Y}} \leq ((2k)^q + 2k) \theta \|x\|_{\mathbf{X}}^q \quad (4.3)$$

for all $x \in \mathbf{X}$. So

$$\left\| f(x) - 2kf\left(\frac{x}{2k}\right) \right\| \leq \frac{(2k)^q + 2k}{(2k)^q - 2k} \theta \|x\|_{\mathbf{X}}^q \quad (4.4)$$

Hence we have

$$\begin{aligned} & \left\| (2k)^l f\left(\frac{x}{(2k)^l}\right) - (2k)^m f\left(\frac{x}{(2k)^m}\right) \right\|_{\mathbf{Y}} \\ & \leq \sum_{j=l}^{m-1} \left\| (2k)^j f\left(\frac{x}{(2k)^j}\right) - (2k)^{j+1} f\left(\frac{x}{(2k)^{j+1}}\right) \right\|_{\mathbf{Y}} \\ & \leq \frac{(2k)^q + 2k}{(2k)^q - 2k} \theta \sum_{j=l}^{m-1} \frac{(2k)^j}{(2k)^{qj}} \theta \|x\|_{\mathbf{X}}^q. \end{aligned} \quad (4.5)$$

for all nongnegative m and l with $m > l$, $\forall x \in \mathbf{X}$. It follows from (4.5) that the sequence $\{(2k)^n f\left(\frac{x}{(2k)^n}\right)\}$ is a cauchy sequence for all $x \in \mathbf{X}$. Since \mathbf{Y} is a Banach space, the

sequence $\{(2k)^n f\left(\frac{x}{(2k)^n}\right)\}$ converges.

So one can define the mapping $H : \mathbf{X} \rightarrow \mathbf{Y}$ by

$$H(x) := \lim_{n \rightarrow \infty} (2k)^n f\left(\frac{x}{(2k)^n}\right)$$

for all $x \in \mathbf{X}$.

$$\begin{aligned} & \left\| \sum_{j=1}^k H(x_j) + \sum_{j=1}^k H(y_j) + \sum_{j=1}^k H(z_j) \right\|_{\mathbf{Y}} \\ &= \lim_{n \rightarrow \infty} (2k)^n \left\| \sum_{j=1}^k f\left(\frac{x_j}{(2k)^n}\right) + \sum_{j=1}^k f\left(\frac{y_j}{(2k)^n}\right) + \sum_{j=1}^k f\left(\frac{z_j}{(2k)^n}\right) \right\|_{\mathbf{Y}} \\ &\leq \lim_{n \rightarrow \infty} (2k)^n \left\| 2kf\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{(2k)^{n+1}}\right) \right\|_{\mathbf{Y}} \\ &\quad + \lim_{n \rightarrow \infty} \frac{(2k)^n}{(2k)^{kq}} \theta \left(\sum_{j=1}^k \|x_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|y_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|z_j\|_{\mathbf{X}}^q \right) \\ &= \left\| 2kH\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k}\right) \right\|_{\mathbf{Y}} \end{aligned} \tag{4.6}$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow k$. So

$$\begin{aligned} & \left\| \sum_{j=1}^k H(x_j) + \sum_{j=1}^k H(y_j) + \sum_{j=1}^k H(z_j) \right\|_{\mathbf{Y}} \\ &\leq \left\| 2kH\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k}\right) \right\|_{\mathbf{Y}} \end{aligned} \tag{4.7}$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$. By Proposition 3.1, the mapping $H : \mathbf{X} \rightarrow \mathbf{Y}$ is additive. Now, let $T : \mathbf{X} \rightarrow \mathbf{Y}$ be another additive mapping satisfy (4.2) then we have

$$\begin{aligned} \|H(x) - T(x)\|_{\mathbf{Y}} &= (2k)^n \left\| h\left(\frac{x}{(2k)^n}\right) - T\left(\frac{x}{(2k)^n}\right) \right\|_{\mathbf{Y}} \\ &\leq (2k)^n \left(\left\| h\left(\frac{x}{(2k)^n}\right) - f\left(\frac{x}{(2k)^n}\right) \right\|_{\mathbf{Y}} + \left\| T\left(\frac{x}{(2k)^n}\right) - f\left(\frac{x}{(2k)^n}\right) \right\|_{\mathbf{Y}} \right) \\ &\leq \frac{2((2k)^n + 2k)}{(2k)^n - 2k} \cdot \frac{\theta}{k} \cdot \frac{(2k)^n}{(2k)^{nq}} \|x\|_{\mathbf{X}}^q. \end{aligned} \tag{4.8}$$

which tends to zero as $q \rightarrow \infty$ for all $x \in \mathbf{X}$. So we can conclude that $H(x) = T(x)$ for all $x \in \mathbf{X}$. This proves the uniqueness of H . Thus the mapping $H : \mathbf{X} \rightarrow \mathbf{Y}$ is additive mapping satisfying (4.2).

Theorem 4.2. Suppose $q < 1$, θ be positive real numbers and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned}
 & \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} \\
 & \leq \left\| 2kf \left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k} \right) \right\|_{\mathbf{Y}} \\
 & + \theta \left(\sum_{j=1}^k \|x_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|y_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|z_j\|_{\mathbf{X}}^q \right) \quad (4.9)
 \end{aligned}$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbf{Y}} \leq \frac{2k + (2k)^q}{2k - (2k)^q} \theta \|x\|_{\mathbf{X}}^q. \quad (4.10)$$

for all $x \in \mathbf{X}$.

Notice that: Form (4.3) we have

$$\left\| f(x) - \frac{1}{2k} f(2kx) \right\|_{\mathbf{Y}} \leq \frac{2k + (2k)^q}{2k} \theta \|x\|_{\mathbf{X}}^q \quad (4.11)$$

for all $x \in \mathbf{X}$. The rest of the proof is similar to the proof of Theorem 4.1.

Theorem 4.3. Suppose $q > p^{-1}$ with $p \geq 3$, θ be non-negative real and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned}
 & \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} \\
 & \leq \left\| 2kf \left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k} \right) \right\|_{\mathbf{Y}} \\
 & + \theta \prod_{j=1}^k \|x_j\|_{\mathbf{X}}^q \cdot \prod_{j=1}^k \|y_j\|_{\mathbf{X}}^q \cdot \|z_1\|_{\mathbf{X}}^{kq} \cdot \left(1 + \prod_{j=2}^k \|z_j\|_{\mathbf{X}}^q \right) \quad (4.12)
 \end{aligned}$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbf{Y}} \leq \frac{(2k)^q}{(2k)^{3kq} - 2k} \theta \|x\|_{\mathbf{X}}^{3kq}. \quad (4.13)$$

for all $x \in \mathbf{X}$.

Proof. Assume that $f: \mathbf{X} \rightarrow \mathbf{Y}$ satisfies (4.12).

We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $(x, \dots, x, x, \dots, x, -2kx, 0, \dots, 0)$ in (4.12), we have

$$\left\| 2kf(x) - f(2kx) \right\|_{\mathbf{Y}} = \left\| 2kf(x) + f(-2kx) \right\|_{\mathbf{Y}} \leq |2k|^{kq} \theta \|x\|_{\mathbf{X}}^{3kq} \quad (4.14)$$

for all $x \in \mathbf{X}$. So

$$\left\| f(x) - 2kf\left(\frac{x}{2k}\right) \right\| \leq \frac{|2k|^{kq}}{|2k|^{3kq}} \|x\|_{\mathbf{X}}^{3kq} \quad (4.15)$$

Hence we have

$$\begin{aligned}
 & \left\| (2k)^l f\left(\frac{x}{(2k)^l}\right) - (2k)^m f\left(\frac{x}{(2k)^m}\right) \right\|_{\mathbf{Y}} \\
 & \leq \sum_{j=l}^{m-1} \left\| (2k)^j f\left(\frac{x}{(2k)^j}\right) - (2k)^{j+1} f\left(\frac{x}{(2k)^{j+1}}\right) \right\|_{\mathbf{Y}} \\
 & \leq \frac{|2k|^{kq}}{|2k|^{3kq}} \sum_{j=l}^{m-1} \frac{(2k)^j}{(2k)^{3kqj}} \theta \|x\|_{\mathbf{X}}^{3kq}. \tag{4.16}
 \end{aligned}$$

for all nongnegative m and l with $m > l$, $\forall x \in \mathbf{X}$. It follows from (4.16) that the sequence

$\left\{ (2k)^n f\left(\frac{x}{(2k)^n}\right) \right\}$ is a cauchy sequence for all $x \in \mathbf{X}$. Since \mathbf{Y} is a Banach space, the sequence $\left\{ (2k)^n f\left(\frac{x}{(2k)^n}\right) \right\}$ converges.

So one can define the mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ by

$$H(x) := \lim_{n \rightarrow \infty} (2k)^n f\left(\frac{x}{(2k)^n}\right)$$

for all $x \in \mathbf{X}$. Moreover, letting $l = 0$ and passing the limit $m \rightarrow \infty$ in (4.16), we have (4.13). The rest of the Prooft is similar to the Proof of the Theorem 4.1.

Theorem 4.4. Suppose $q < p^{-1}$ with $p \geq 3$, θ be non-negative real and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned}
 & \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} \\
 & \leq \left\| 2k f\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k}\right) \right\|_{\mathbf{Y}} \\
 & \quad + \theta \prod_{j=1}^k \|x_j\|_{\mathbf{X}}^q \cdot \prod_{j=1}^k \|y_j\|_{\mathbf{X}}^q \cdot \|z_1\|_{\mathbf{X}}^{kq} \cdot \left(1 + \prod_{j=2}^k \|z_j\|_{\mathbf{X}}^q\right) \tag{4.17}
 \end{aligned}$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\|f(x) - H(x)\|_{\mathbf{Y}} \leq \frac{(2k)^q}{2k - (2k)^{3kq}} \theta \|x\|_{\mathbf{X}}^{3kq}. \tag{4.18}$$

for all $x \in \mathbf{X}$.

The rest of the Prooft is similar to the Proof of the Theorem 4.1.

5. Establishing solutions to functional inequality (1.2) related to the type of Cauchy additive functional equation

Now, we first study the solutions of (1.2). Note that for this inequality, \mathbf{X} is a normed space with norm $\|\cdot\|_{\mathbf{X}}$ and that \mathbf{Y} is a Banach space with norm $\|\cdot\|_{\mathbf{Y}}$. Under this setting, we can show that the mappings satisfying (1.2) is Cauchy additive. These results are give in the following.

Theorem 5.1. Suppose $q > 1$, θ be non-negative real and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned}
 & \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} \leq \left\| f\left(\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j\right) \right\|_{\mathbf{Y}} \\
 & \quad + \theta \left(\sum_{j=1}^k \|x_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|y_j\|_{\mathbf{X}}^q + \sum_{j=1}^k \|z_j\|_{\mathbf{X}}^q \right) \tag{5.1}
 \end{aligned}$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbf{Y}} \leq \frac{(2k)^q + 2k}{(2k)^q - 2k} \theta \left\| x \right\|_{\mathbf{X}}^q. \quad (5.2)$$

for all $x \in \mathbf{X}$.

The rest of the Proof is similar to the Proof of the Theorem 4.1.

Theorem 5.2. Suppose $q < 1$, θ be positive real numbers and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be a mapping such that

$$\begin{aligned} \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} &\leq \left\| f\left(\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j\right) \right\|_{\mathbf{Y}} \\ &+ \theta \left(\sum_{j=1}^k \left\| x_j \right\|_{\mathbf{X}}^q + \sum_{j=1}^k \left\| y_j \right\|_{\mathbf{X}}^q + \sum_{j=1}^k \left\| z_j \right\|_{\mathbf{X}}^q \right) \end{aligned} \quad (5.3)$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\left\| \frac{f(x) + f(-x)}{2k} - H(x) \right\|_{\mathbf{Y}} \leq \frac{2k + (2k)^q}{2k - (2k)^q} \theta \left\| x \right\|_{\mathbf{X}}^q. \quad (5.4)$$

for all $x \in \mathbf{X}$.

The rest of the proof is similar to the proof of Theorem 4.1 and 4.2.

Theorem 5.3. Suppose $q > p^{-1}$ with $p \geq 3$, θ be non-negative real and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned} \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} &\leq \left\| f\left(\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j\right) \right\|_{\mathbf{Y}} \\ &+ \theta \prod_{j=1}^k \left\| x_j \right\|_{\mathbf{X}}^q \cdot \prod_{j=1}^k \left\| y_j \right\|_{\mathbf{X}}^q \cdot \left\| z_1 \right\|_{\mathbf{X}}^{kq} \cdot \left(1 + \prod_{j=2}^k \left\| z_j \right\|_{\mathbf{X}}^q \right) \end{aligned} \quad (5.5)$$

for all $x_j, y_j, z_j \in \mathbf{X}$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: \mathbf{X} \rightarrow \mathbf{Y}$ such that

$$\left\| f(x) - H(x) \right\|_{\mathbf{Y}} \leq \frac{(2k)^q}{(2k)^{3kq} - 2k} \theta \left\| x \right\|_{\mathbf{X}}^{3kq}. \quad (5.6)$$

for all $x \in \mathbf{X}$.

Proof. Assume that $f: \mathbf{X} \rightarrow \mathbf{Y}$ satisfies (5.5).

We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $(x, \dots, x, x, \dots, x, -2kx, 0, \dots, 0)$ in (5.5), we have

$$\left\| 2kf(x) - f(2kx) \right\|_{\mathbf{Y}} = \left\| 2kf(x) + f(-2kx) \right\|_{\mathbf{Y}} \leq |2k|^{kq} \theta \left\| x \right\|_{\mathbf{X}}^{3kq} \quad (5.7)$$

for all $x \in \mathbf{X}$. The rest of the Proof is similar to the Proof of the Theorem 4.1 and 4.3.

Theorem 5.4. Suppose $q < p^{-1}$ with $p \geq 3$, θ be non-negative real and $f: \mathbf{X} \rightarrow \mathbf{Y}$ be an odd mapping such that

$$\begin{aligned} \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_{\mathbf{Y}} &\leq \left\| 2kf\left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j + \sum_{j=1}^k z_j}{2k}\right) \right\|_{\mathbf{Y}} \\ &+ \theta \prod_{j=1}^k \left\| x_j \right\|_{\mathbf{X}}^q \cdot \prod_{j=1}^k \left\| y_j \right\|_{\mathbf{X}}^q \cdot \left\| z_1 \right\|_{\mathbf{X}}^{kq} \cdot \left(1 + \prod_{j=2}^k \left\| z_j \right\|_{\mathbf{X}}^q \right) \end{aligned} \quad (5.8)$$

for all $x_j, y_j, z_j \in X$ for all $j = 1 \rightarrow n$

. Then there exists a unique additive mapping $H: X \rightarrow Y$ such that

$$\|f(x) - H(x)\|_Y \leq \frac{(2k)^q}{2k - (2k)^{3kq}} \theta \|x\|_X^{3kq}. \quad (5.9)$$

for all $x \in X$.

The rest of the Proof is similar to the Proof of the Theorem 4.1 and 4.4.

6. Establishing solutions to functional inequality (1.3) related to the type of Cauchy-Jensen additive functional equation
 Now, we first study the solutions of (1.3). Note that for this inequality, X is a normed space with norm $\|\cdot\|_X$ and that Y is a Banach space with norm $\|\cdot\|_Y$. Under this setting, we can show that the mappings satisfying (1.3) is Cauchy-Jensen additive. These results are give in the following.

Theorem 6.1. Suppose $q > 1$, θ be non-negative real, $f(0) = 0$ and $f: X \rightarrow Y$ be a mapping such that

$$\begin{aligned} \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + 2k \sum_{j=1}^k f(z_j) \right\|_Y &\leq \left\| 2kf \left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j}{2k} + \sum_{j=1}^k z_j \right) \right\|_Y \\ &\quad + \theta \left(\sum_{j=1}^k \|x_j\|_X^q + \sum_{j=1}^k \|y_j\|_X^q + \sum_{j=1}^k \|z_j\|_X^q \right) \end{aligned} \quad (6.1)$$

for all $x_j, y_j, z_j \in X$ for all $j = 1 \rightarrow n$. Then there exists a unique additive mapping $H: X \rightarrow Y$ such that

$$\|f(x) - H(x)\|_Y \leq \frac{(2k)^q + 1}{(2k)^q - 2k} \theta \|x\|_X^q. \quad (6.2)$$

for all $x \in X$.

Proof. Assume that $f: X \rightarrow Y$ satisfies (6.1).

We replacing $(x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_k)$ by $(2kx, 0, \dots, 0, 0, \dots, 0, -x, 0, \dots, 0)$ in (6.1), we have

$$\left\| f(2kx) - 2kf(x) \right\|_Y = \left\| f(2kx) + 2kf(-x) \right\|_Y \leq ((2k)^q + 1) \theta \|x\|_X^q \quad (6.3)$$

for all $x \in X$. So

$$\left\| f(x) - 2kf\left(\frac{x}{2k}\right) \right\|_Y \leq \frac{(2k)^q + 1}{(2k)^q} \theta \|x\|_X^q \quad (6.4)$$

The rest of the proof is similar to the proof of Theorem 4.1.

Theorem 6.2. Suppose $q < 1$, θ be positive real numbers and $f: X \rightarrow Y$ be a mapping such that

$$\begin{aligned} \left\| \sum_{j=1}^k f(x_j) + \sum_{j=1}^k f(y_j) + \sum_{j=1}^k f(z_j) \right\|_Y &\leq \left\| 2kf \left(\frac{\sum_{j=1}^k x_j + \sum_{j=1}^k y_j}{2k} + \sum_{j=1}^k z_j \right) \right\|_Y \\ &\quad + \theta \left(\sum_{j=1}^k \|x_j\|_X^q + \sum_{j=1}^k \|y_j\|_X^q + \sum_{j=1}^k \|z_j\|_X^q \right) \end{aligned} \quad (6.5)$$

for all $x_j, y_j, z_j \in X$ for all $j = 1 \rightarrow n$

$$\|f(x) - H(x)\|_Y \leq \frac{2k + (2k)^q}{2k - (2k)^q} \theta \|x\|_X^q. \quad (6.6)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 4.1 and 4.2.

5. CONCLUSION

In this paper I have given three general functional inequalities and I have shown that their solutions are determined on normalized spaces and take values in Banach spaces.

REFERENCES

- [1] S.M. ULam A collection of Mathematical problems, volume 8, Interscience Publishers. New York,1960.
- [2] Donald H. Hyers, On the stability of the functional equation, Proceedings of the National Academy of the United States of America, 27(4)(1941),222.<https://doi.org/10.1073/pnas.27.4.222>,
- [3] Th. M. Rassias, On the stability of the linear mapping in Banach spaces, Proceedings of the American Mathematical Society, vol.72,no.2,pp.297-300,1978.
- [4] Z. Gajda, On stability of additive mappings, International Journal of Mathematics and Mathematical Sciences, vol.14,no.3,pp.431-434,1991.
- [5] S. Czerwinski, Functional Equations and Inequalities in Several Variables, World Scientific, River Edge, NJ, USA, 2002.
- [6] P. Găvrută, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," Journal of Mathematical Analysis and Applications, vol.184,no.3,pp.431-436,1994.
- [7] A. Gilănyi, Eine zur Parallelogrammgleichung äquivalente Ungleichung, Aequationes Mathematicae, vol.62,no.3,pp.303-309,2001.
- [8] Th. M. Rassias and P. Semrl, On the behavior of mappings which do not satisfy Hyers-Ulam stability, Proceedings of the American Mathematical Society, vol.114,no.4,pp.989-993,1992.
- [9] D. H. Hyers, G. Isac, and Th. M. Rassias, Stability of Functional Equations in Several Variables, vol. 34 of Progress in Nonlinear Differential Equations and Their Applications, Birkhauser Boston, Boston, Mass, USA, 1998.
- [10] J. M. Rassias, On approximation of approximately linear mappings by linear mappings, Journal of Functional Analysis, vol.46,no.1,pp.126-130,1982.
- [11] K.-W. Jun and Y.-H. Lee, A generalization of the Hyers-Ulam-Rassias stability of the pexiderized quadratic equations, Journal of Mathematical Analysis and Applications, vol.297,no.1,pp.7086, 2004.
- [12] S.-M. Jung, Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis, Hadronic Press, Palm Harbor, Fla, USA, 2001.
- [13] C. Park, Homomorphisms between Poisson JC-algebras, Bulletin of the Brazilian Mathematical Society. New Series, vol.36,no.1,pp.79-97,2005.
- [14] C. Park, Hyers-Ulam-Rassias stability of homomorphisms in quasi-Banach algebras to appear in Bulletin des Sciences Mathématiques.
- [15] Th. M. Rassias, Problem 16; 2, Report of the 27th International Symp. on Functional Equations, Aequationes Mathematicae, vol.39,pp.292-93;309,1990
- [16] J. Ratz, On inequalities associated with the Jordan-von Neumann functional equation, Aequationes Mathematicae, vol.66,no.1-2,pp.191-200,2003.ChoonkilParketal.13
- [17] A. Gilănyi, On a problem by K. Nikodem,Mathematical Inequalities Applications, vol. 5, no. 4, pp. 707-710, 2002.
- [18] T. Aoki, On the stability of the linear transformation in Banach space, J. Math. Soc. Japan 2(1950), 64-66.
- [19] A. Bahyrycz, M. Piszczek, Hyers stability of the Jensen function equation, Acta Math. Hungar.,142 (2014),353-365.
- [20] M. Balcerowski, On the functional equations related to a problem of Z. Boros and Z. Dro'czy, Acta Math. Hungar.,138 (2013), 329-340.
- [21] W. Fechner, Stability of a functional inequalities associated with the Jordan-von Neumann functional equation, Aequationes Math. 71 (2006), 149-161.
- [22] W. P and J. Schwaiger, A system of two inhomogeneous linear functional equations, Acta Math. Hungar 140 (2013), 377-406.
- [23] L. Maligranda. Tosio Aoki (1910-1989) in International symposium on Banach and function spaces:14/09/2006-17/09/2006, pages 1-23. Yokohama Publishers, 2008.
- [24] A. Najati and G.Z. Eskandani. Stability of a mixed additive and cubic functional equation in quasi- Banach spaces. J. Math. Anal. Appl.342(2):1318–1331, 2008.
- [25] Attila Gilănyi, On a problem by K. Nikodem, Math. Inequal. Appl., 5 (2002), 707-710.
- [26] W. Fechner, On some functional inequalities related to the logarithmic mean, Acta Math., Hungar., 128 (2010),31-45, 303-309.
- [27] Choonkil.Park. Additive β -functional inequalities, Journal of Nonlinear Science and Appl. 7(2014), 296-310.
- [28] LY VAN AN, Hyers-Ulam stability of functional inequalities with three variable in Banach spaces and Non-Archimedean Banach spaces International Journal of Mathematical Analysis Vol.13, 2019, no. 11. 519-53014, 296-310 . <https://doi.org/10.12988/ijma.2019.9954>.
- [29] Choonkil Park, Functional in equalities in Non-Archimedean normed spaces. Acta Mathematica Sinica, English Series, 31 (3), (2015), 353-366 <https://doi.org/10.1007/s10114-015-4278-5>.
- [30] Jung Rye Lee, Choonkil* Park, and Dong Yun Shin Additive and quadratic functional in equalities in Non-Archimedean normed spaces, International Journal of Mathematical Analysis, 8 (2014), 1233- 1247.

- [31] Y.J.Cho^a, Choonkil Park^{b*}, and R.Saadati^{c,b*} functional in equalities in Non-Archimedean normed spaces, *Applied Mathematics Letters* 23(2010), 1238-1242.
- [32] Y.Aribou*, S.Kabbaj Generalized functional in inequalities in Non-Archimedean normed spaces, *Applied Mathematics Letters* 2, (2018) Pages: 61-66.
- [33] LY VAN AN, Hyers-Ulam stability additive β -functional inequalities with three variable in Banach spaces and Non-Archimedean Banach spaces *International Journal of Mathematical Analysis* Vol.14, 2020, no. 5-8. 519-53014, 296-310 .<https://doi.org/10.12988/ijma.2020.91169>.
- [34] LY VAN AN,Generalized Hyes-Ulam stability of the additive functional inequalities with 2n- variables in non-Archimedean Banach space. *Bulletin of mathematics and statistics research.* Vol.9.Issue.3.2021 (July-Sept) Ky publications .<http://www.bomsr.comDOI:10.33329/bomsr.9.3. 67>.
- [35] , Qarawani, M.N. (2013), Hyers-Ulam-Rassias Stability for the Heat Equation, *Applied Mathematics*, Vol. 4 No. 7, 2013, pp. 1001-1008. doi:10.4236/am.2013.47137.