

Computing best discrete least-squares approximations by first-degree splines with free knots

Ludwig J. Cromme¹, Jens Kunath² and Andreas Krebs³

Abstract

We present an algorithm to compute best least-squares approximations of discrete real-valued functions by first-degree splines (broken lines) with free knots. We demonstrate that the algorithm delivers after a finite number of steps a (global) best approximation. The analysis is complemented by remarks on programming and by a number of numerical examples including applications from medicine (MBC, MIC).

Key words: splines, first-degree splines, broken lines, splines with free knots, least-squares approximation, best approximation, minimal bactericidal concentration (MBC), minimal inhibitory concentration (MIC)

1. Introduction

Splines of a real variable with free knots form a highly nonlinear approximation family with varying dimension of the boundary (Cromme [2]). Even just local best approximations are therefore not easily computed (Cromme [3]) and globally convergent numerical methods are not known.

But for the special case of first-degree splines (broken lines) with free knots a globally convergent numerical method for the approximation of discrete data in the least-squares sense could be developed and is presented in the following sections.

The basic idea of the method is to reduce the complexity of the original problem: Instead of solving the original nonlinear problem on the whole data set a number of just linear problems on subsets ("segments") is treated and assembled to a solution of the original problem.

The underlying idea is sketched in Fig. 1: In this figure the best L_2 -approximation s^* from the set of first-degree splines with 5 free knots t_1, \dots, t_5 to discrete data (x_i, f_i) , $i = 0, 1, \dots, 18$, is plotted. Knots can be located between two data abscissae (such knots will be called interior knots in the following, in our example t_1 and t_4 are interior knots) or coincide with data abscissae (called data knots, in our example $t_2 = x_4, t_3 = x_{10}$, and $t_5 = x_{15}$ are data knots).

Basic is the observation that s^* is not only best approximation on the whole data set, but also on subsets of the data called segments; for a formal introduction of notations see below. For example $s^*|_{[x_4, x_{12}]}$ is a best approximation from $S^1(x_7, x_{10})$, the first-degree splines with fixed knots x_7 and x_{10} , to the data $(x_4, f_4), \dots, (x_{12}, f_{12})$. For if a better approximation \tilde{s} with fixed knots x_7 and x_{10} to the data $(x_4, f_4), \dots, (x_{12}, f_{12})$ existed, s^* could be modified in the considered segment in the direction of \tilde{s} . The interior knots t_1 and t_4 would be slightly moved in this process, the other knots would remain unchanged. The approximation error of the thus modified spline would be smaller than the approximation error of s^* - contradicting the optimality of s^* as best approximation.

So if these segments ranging from one interior knot to the next interior knot were known, we could determine s^* by solving just *linear (!)* approximation problems with fixed knots segment for segment. But since the segments of a best approximation are not known in advance our algorithm examines systematically possible segmentations of the data abscissae to see if they might lead to a best approximation.

¹Numerische und Angewandte Mathematik, BTU Cottbus-Senftenberg, Platz der Deutschen Einheit 1, D-03046 Cottbus, Germany, E-mail: Ludwig.Cromme@b-tu.de

²Numerische und Angewandte Mathematik, BTU Cottbus-Senftenberg, Platz der Deutschen Einheit 1, D-03046 Cottbus, Germany

³Aerodynamik und Strömungslehre, BTU Cottbus-Senftenberg, Siemens-Halske-Ring 14, D-03046 Cottbus, Germany




Figure 1: A first-degree spline (broken line) $s^* \in S_5^1[a, b]$ approximating the $\mu + 2 = 20$ data $(x_0, f_0), \dots, (x_{19}, f_{19})$. s^* has $k = 5$ simple knots t_1, \dots, t_5 and the boundary knots $t_0 := a := x_0$ and $t_6 := b := x_{19}$.

The implementation of this idea raises a number of questions which will be addressed after formal presentation of the algorithm in the following section.

Let us summarize the most important notations used: Let \mathbb{P}_m denote the real-valued polynomials of degree smaller or equal m and $[a, b]$ a real interval $a < b$. For $a =: t_0 < t_1 < \dots, < t_k < t_{k+1} := b$ and $m, k \in \mathbb{N}$ let

$$S^m(t_1, \dots, t_k) := \{s \in C^{m-1}[a, b] \mid s_j|_{(t_j, t_{j+1})} \in \mathbb{P}_m, j = 0, 1, \dots, k\}$$

be the set of splines of degree m with k fixed (simple) knots t_1, \dots, t_k . The splines of degree $m \in \mathbb{N}$ with at most $k \in \mathbb{N}$ free (simple) knots are the set

$$S_k^m[a, b] := \left\{ s \in C^{m-1}[a, b] \mid \begin{array}{l} \text{there are points} \\ a =: t_0 < t_1 < \dots < t_k < t_{k+1} := b, \\ \text{such that } s|_{(t_i, t_{i+1})} \in \mathbb{P}_m \text{ for } i = 0, 1, \dots, k \end{array} \right\} .$$

t_i is called an *improper* (or *inactive*) knot of a spline $s \in S_k^m[a, b]$ if s is m -times differentiable in t_i , otherwise the knot is called *proper* (or *active*) knot.

Let $\mu + 2$ real abscissae $a =: x_0 < x_1 < \dots < x_{\mu+1} := b$ and function values $f_0, \dots, f_{\mu+1} \in \mathbb{R}$ of a function $f : [a, b] \rightarrow \mathbb{R}$ be given. We define vectors $X := (x_0, \dots, x_{\mu+1})^t$, $F := (f_0, \dots, f_{\mu+1})^t$, and for $g : \{x_0, \dots, x_{\mu+1}\} \rightarrow \mathbb{R}$ the vector $g(X) := (g(x_0), \dots, g(x_{\mu+1}))^t$. With the Euclidian norm $\|\cdot\|_2$ our problem can now be stated:

Definition 1.1. (Approximation problem) Find $s^* \in S_k^1[a, b]$ with

$$\|F - s^*\|_{2, X} := \|F - s^*(X)\|_2 = \inf_{s \in S_k^1[a, b]} \|F - s(X)\|_2 .$$

In the following sections we generally assume $k \geq 1$ and $\mu \geq k + 1 \geq 2$, since otherwise the problem would be a purely polynomial approximation problem or the data could be reproduced exactly in $S_k^1[a, b]$.

2. The basic algorithm and convergence theorem

A spline $s \in S_k^1[a, b]$ partitions the data abscissae $x_0 < x_1 < \dots < x_{\mu+1}$ naturally into segments each ranging from an interior knot (i.e. a knot situated between two data abscissae) to the next interior knot. Within a segment there can be more data knots, i.e. knots coinciding with data abscissae.

Such segmentations are basic for the numerical procedure presented below. In defining the segments it is advantageous to code the position of knots in a "position vector", i.e. the information whether a knot coincides with an abscissa and with which or between which abscissae a knot is located. To start we define position vectors for arbitrary first-degree splines. As will be seen later, the algorithm needs to deal only with "regular" position vectors, see Definition 2.6.

Definition 2.1. (a) A position vector p of length k to the data $x_0, \dots, x_{\mu+1}$ is a vector from \mathbb{N}_0^k with

$$0 \leq p(j) \leq p(j+1) \leq 2\mu, \quad j = 1, 2, \dots, k-1,$$

$$p(j) = p(j+1) \Rightarrow p(j) \text{ is even,}$$

where $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ and 0 is considered even.

(b) For $s \in S_k^1[a, b]$, $a =: x_0 < x_1 < \dots < x_{\mu+1} := b$ and $x_0 < t_1 < t_2 < \dots < t_k < x_{\mu+1}$ the position vector $p \in \mathbb{N}_0^k$ of s to the data $x_0, \dots, x_{\mu+1}$ is defined as

$$p(j) := \begin{cases} 2i-1 & , \text{ if } t_j = x_i \\ 2i & , \text{ if } t_j \in (x_i, x_{i+1}) \end{cases} .$$

Conclusion 2.2. (a) The position vector for s^* from Fig. 1 is the vector $p = (6, 13, 19, 24, 29)$.

(b) A position vector of s allows us to infer back the position of the knots of s : $t_j = x_{\frac{p(j)+1}{2}}$, if $p(j)$ is odd and $t_j \in (x_{\frac{p(j)}{2}}, x_{\frac{p(j)}{2}+1})$, if $p(j)$ is even. By Definition 2.1 (b) the position of knots is mapped one-to-one onto the set of position vectors.

(c) Successive components $p(j)$ and $p(j+1)$ of a position vector p for $s \in S_k^1[a, b]$ are equal iff t_j and t_{j+1} lie in the same interval (x_i, x_{i+1}) :

$$p(j) = p(j+1) \Leftrightarrow x_{\frac{p(j)}{2}} < t_j < t_{j+1} < x_{\frac{p(j)}{2}+1} .$$

(d) For every position vector p according to Definition 2.1 (a) exists a spline $s \in S_k^1[a, b]$ with p as position vector. To be more precise, there exists a whole class of splines all of which have p as corresponding position vector.

Proof. We confine ourselves to the proof of (d). Let p be a position vector according to Definition 2.1 (a). For odd $p(j)$ set $t_j := x_{\frac{p(j)+1}{2}}$. For even $p(j)$ without loss of generality let $p(j)$ to $p(j+q)$ be exactly the components of p identical with $p(j)$, that is, $p(j) = p(j+1) = \dots = p(j+q)$ and $p(j-1) < p(j)$, if $j \geq 2$ and $p(j+q) < p(j+q+1)$, if $j+q < k$. Choose t_j, \dots, t_{j+q} such that

$$x_{\frac{p(j)}{2}} < t_j < t_{j+1} < \dots < t_{j+q} < x_{\frac{p(j)}{2}+1}$$

holds. This implies $t_1 < t_2 < \dots < t_k$ and each spline from $S_k^1[a, b]$ with these knots has position vector p . \square

The position of the knots (coded by the position vector) induces a segmentation of the data. Each segment ranges from one interior knot to the next, more precisely:

Definition/Conclusion 2.3. Let $p \in \mathbb{N}_0^k$ be a position vector for $s \in S_k^1[a, b]$ according to Definition 2.1:

- **Case 1:** All $p(j)$ are odd ($j = 1, 2, \dots, k$). Then s has no interior knots, all knots coalesce with date abscissae and we have

$$t_j := x_{\frac{p(j)+1}{2}} \quad \text{for } j = 1, 2, \dots, k .$$

There is only one segment $S_0 := \{x_0, x_1, \dots, x_{\mu+1}\}$.

- **Case 2:** At least one component $p(j)$ of p is even. Let $r \in \{1, \dots, k\}$ be chosen maximal such that $p(j_l)$ is even for $l = 1, \dots, r$ and suitable j_l with $1 \leq j_1 < j_2 < \dots < j_r \leq k$. The knots t_{j_1}, \dots, t_{j_r} are exactly the interior knots of s . Set

$$i_l := \frac{p(j_l)}{2}, \quad l = 1, 2, \dots, r .$$

Let $r \geq 2$, $l \in \{1, \dots, r-1\}$ hold. In case $i_l = i_{l+1}$, set $S_l := \emptyset$. In case $i_l + 1 \leq i_{l+1}$ set

$$S_l := \{x_{i_l+1}, x_{i_l+2}, \dots, x_{i_{l+1}}\} .$$

If $i_l + 1 \leq i_{l+1}$ and in addition $q_l := j_{l+1} - j_l - 1 > 0$ holds then

$$t_{j_l+i} = x_{\frac{p(j_l+i)+1}{2}} \quad , \quad i = 1, 2, \dots, q_l,$$

are the knots in segment S_l coalescing with data abscissae. So the segments S_0 (to the left of first interior knot t_{j_1}) and S_r (to the right of the last interior knot t_{j_r}) are left to be defined here (and also for $r = 1$). S_0 consists of abscissae x_0, \dots, x_{i_1} and contains - if $j_1 - 1 > 0$ - the knots

$$t_i = x_{\frac{p(i)+1}{2}} \quad , \quad i = 1, 2, \dots, j_1 - 1 \quad .$$

Segment S_r consists of abscissae $x_{i_r+1}, \dots, x_{\mu+1}$ and contains - if $j_r + 1 \leq k$ - the knots

$$t_i = x_{\frac{p(i)+1}{2}} \quad , \quad i = j_r + 1, \dots, k \quad .$$




Figure 2: Diagram of segment S_l : Numbers $0, 1, 2, \dots, 2\mu$ as pointer for the position of knots and the segment $S_l = \{x_{i_l+1}, \dots, x_{i_{l+1}}\}$ ranging from interior knot t_{j_l} to interior knot $t_{j_{l+1}}$.

Example 2.4. Position vector $p = (6, 1, 3, 19, 24, 29)$ for spline s^* in Fig. 1 leads to segments $S_0 = \{x_0, x_1, x_2, x_3\}$ without knots, $S_1 = \{x_4, x_5, \dots, x_{12}\}$ with knots $t_2 := x_7$ and $t_3 := x_{10}$ and to segment $S_2 = \{x_{13}, x_{14}, \dots, x_{18}\}$ with the knot $t_5 := x_{15}$.

Remark 2.5. Because of Theorem 12 in [4] we can limit our search for a best approximation to splines with only simple knots and for which the additional features (a) to (d) from Theorem 12 in [4] hold; see also Appendix A. In Lemma 2.7 we will prove that these are exactly the splines with regular position vectors.

Definition 2.6. A position vector $p \in \mathbb{N}_0^k$ is called *regular position vector of length k*, if the following features hold:

- (a) $\{p(1), \dots, p(k)\} \subseteq \{1, 2, \dots, 2\mu - 1\}$ and
- (b) $p(j)$ even $\Rightarrow \forall i \in \{1, \dots, j - 1\} : p(i) \neq p(j) - 1, \forall i \in \{j + 1, \dots, k\} : p(i) \neq p(j) + 1$.
- (c) For $p(j)$ and $p(j + l + 1)$ ($l \geq 0$) both even and for $l \geq 1$ in addition $p(j + 1), \dots, p(j + l)$ odd the following inequality holds

$$\frac{p(j + l + 1)}{2} - \frac{p(j)}{2} \geq 2 + l \quad .$$

- (d) $\forall j \in \{1, \dots, k - 1\} : p(j + 1) - p(j) \geq \begin{cases} 4 & , \text{ for } p(j), p(j + 1) \text{ both even} \\ 2 & , \text{ otherwise} \end{cases}$

Lemma 2.7. The position vector of a spline $s \in S^k[a, b]$ is regular iff s exhibits features (a) thru (d) from Theorem 12 in [4].

Proof. Lemma 2.7 reflects the fact that properties (a) to (d) in Definition 2.6 for components $p(1), \dots, p(k)$ guarantee features (a) to (d) in Theorem 12 in [4] and vice versa. As an example let us elaborate (d):

So let p be a position vector of a spline $s \in S_k^1[a, b]$, t_j and t_{j+1} knots of s .

- Case 1: t_j and t_{j+1} are interior knots, i.e. $p(j)$ and $p(j+1)$ are even. Then at least two data abscissae are situated between t_j and t_{j+1} iff

$$\frac{p(j+1)}{2} - \left(\frac{p(j)}{2} + 1 \right) + 1 \geq 2 \quad \Leftrightarrow \quad p(j+1) - p(j) \geq 4 \quad .$$

- Case 2: t_j is an interior and t_{j+1} a data knot, i.e., $p(j)$ is even and $p(j+1)$ is odd. Then at least two data abscissae lie between or on t_j and t_{j+1} iff

$$\frac{p(j)}{2} + 1 < \frac{p(j+1) + 1}{2} \Leftrightarrow p(j+1) - p(j) > 1 \Leftrightarrow p(j+1) - p(j) \geq 2 \quad .$$

The equivalence of feature (d) from Theorem 12 in [4] to the requirement

$$p(j+1) - p(j) \geq 2$$

is shown with similar arguments in

- Case 3: t_j is data knot, t_{j+1} interior knot, that is, $p(j)$ is odd and $p(j+1)$ is even,
- Case 4: t_j and t_{j+1} are data knots, that is, $p(j)$ and $p(j+1)$ are odd.

Altogether this proves the equivalence of 2.6 (d) and Theorem 12 (d) from [4]. □

Conclusion 2.8. For a regular position vector the following statements hold true:

$$\{p(1), \dots, p(k)\} \subseteq \{1, 2, \dots, 2\mu - 1\} \quad \text{and} \quad p(1) < p(2) < \dots < p(k) \quad .$$

Proof. See Definition 2.6 (a) and (d).

Lemma 2.9. The number of regular position vectors grows exponentially in μ .

Proof. According to Definition 2.6 a position vector is regular if all k components $p(1) < p(2) < \dots < p(k)$ are odd, i.e.

$$p(j) \in \{1, 3, 5, \dots, 2\mu - 3, 2\mu - 1\}, \quad j = 1, \dots, k \quad .$$

In such cases all knots of related splines are data knots. So there are at least as many regular position vectors as there are subsets of $\{1, 3, 5, \dots, 2\mu - 1\}$ with k elements. According to standard results in combinatorics there are $\binom{\mu}{\mu-k} = \frac{\mu!}{(\mu-k)!k!}$ such subsets. With growing μ and $k = \frac{\mu}{2}$ for example exponential growth of the number of regular position vectors is therefore programmed. □

The fast growth of the number of regular position vectors is evident in Table 1:

We can now specify the algorithm for the solution of Approximation problem 1.1: The algorithm examines systematically all regular position vectors to see if they might code the position of the knots of a best approximating spline $s \in S_k^1[a, b]$.

Algorithm 2.10.

1. Set $\varrho := \infty$.
2. For the next regular position vector p of length k for the data abscissae $x_0, \dots, x_{\mu+1}$ which has not yet been examined carry out exactly one of the following two steps 3 and 4:
3. If all $p(i)$ for $i = 1, 2, \dots, k$ are odd compute the best approximation $s \in S^1 \left(x_{\frac{p(1)-1}{2}}, \dots, x_{\frac{p(k)-1}{2}} \right)$ to the data $(x_0, f_0), \dots, (x_{\mu+1}, f_{\mu+1})$. The spline s is thus the solution of a discrete linear least-squares approximation problem with fixed knots. If $\|F - s(X)\|_2 < \varrho$, set $\varrho := \|F - s(X)\|_2$, $s^* := s$. Continue with step 2.

Table 1: Number of regular position vectors as a function of knots and data points.

number of knots k	number of data points $\mu + 2$					
	15	20	25	30	35	40
1	25	35	45	55	65	75
2	265	545	925	1405	1985	2665
3	1561	4991	11521	22151	37881	59711
4	5641	29961	97281	241601	506921	947241
5	13073	124515	590557	1937199	5060441	11326283
6	19825	369305	2668525	11847485	39146185	106114625
7	19825	795455	9173505	56610575	249673265	799538175

4. If at least one component $p(i)$ is even, i.e., for a suitable $r \in \{1, \dots, k\}$ there are exactly r interior knots, carry out the following steps; notations are as in Definition 2.3, in addition $i_0 := -1, i_{r+1} := \mu + 2, j_0 := 0, j_{r+1} := k + 1, q_0 := j_1 - 1, q_r := k - j_r$:

(a) For each segment S_l ($l = 0, 1, \dots, r$) compute the best least-squares approximation s_l to the data $(x_{i_{l+1}}, f_{i_{l+1}}), \dots, (x_{i_{l+1}+q_l}, f_{i_{l+1}+q_l})$. Here s_l is to chosen from \mathbb{P}_1 if $q_l = 0$ and

$$s_l \in S^1(t_{j_{l+1}}, \dots, t_{j_{l+q_l}}) = S^1\left(x_{\frac{p(j_{l+1})+1}{2}}, \dots, x_{\frac{p(j_{l+1}+q_l)+1}{2}}\right)$$

if $q_l > 0$.

(b) For all $l = 1, \dots, r$ check whether the splines s_{l-1} and s_l intersect in some point $z_l \in (x_{i_l}, x_{i_{l+1}})$. If this is the case for all $l, l = 1, \dots, r$ define

$$s(x) := \begin{cases} s_0(x) & , \quad x_0 \leq x \leq z_1 \\ s_l(x) & , \quad z_l < x \leq z_{l+1}, \quad 1 \leq l \leq r - 1 \\ s_r(x) & , \quad z_r < x \leq x_{\mu+1} \end{cases} .$$

If $\|F - s(X)\|_2 < \varrho$ set $\varrho := \|F - s(X)\|_2, s^* := s$ and procede with step 2. If for some $l \in \{1, \dots, r\}$ splines s_{l-1} and s_l do not intersect in $(x_{i_l}, x_{i_{l+1}})$ reject the position vector p under investigation and continue with step 2.

Procede with step 2.

5. When all regular position vectors p of length k have been examined s^* is the wanted best approximation and ϱ the corresponding approximation error. End.

The proof that Algorithm 2.10 solves the approximation problem is based on Theorem 12 in [4]:

Theorem 2.11. *Algorithm 2.10 stops after a finite number of steps and delivers with s^* a best approximation*

$$\|f - s^*\|_{2,X} = \inf_{s \in S_k^1[x_0, x_{\mu+1}]} \|f - s\|_{2,X} .$$

Proof. Let s^* be a best approximation whith the additional features from Theorem 12 in [4]. Because of Lemma 2.7 the position vector p^* of s^* is then regular and is therefore treated in Algorithm 2.10.

Furthermore, the following assertion holds:

Assertion 2.12. *When Algorithm 2.10 elaborates on p^* the function computed in step 3 or step 4 by the algorithm ist just s^* .*

Proof. a) If step 3 is executed (all $p^*(i)$ are odd) then s^* as best approximation from $S_k^1[a, b]$ is even more best approximation from the smaller set $S^1\left(x_{\frac{p(1)+1}{2}}, \dots, x_{\frac{p(k)+1}{2}}\right)$ which s^* is also an element of.

b) If step 4 is executed every approximation s_i on a segment is identical with s^* in its domain. For simplicity we demonstrate this only for the first segment S_0 .

Assumption: s_0 is a better linear approximation to the data $(x_0, f_0), \dots, \left(x_{\frac{p(j_1)}{2}}, f_{\frac{p(j_1)}{2}}\right)$ than s^* . Any sufficiently small shift of s^* in the direction of s_0 improves the approximation error. Because $(s^*)'$ has a jump discontinuity in t_{j_1} (see Theorem 12 (h) in [4]), the modification of s^* in direction s_0 as described above results in a $s_{new}^* \in S_k^1[a, b]$ with $\|F - s_{new}^*(X)\|_2 < \|F - s^*(X)\|_2$ contradicting the optimality of s^* .

The above assumption must therefore be wrong and the assertion b) holds. s^* is thus an element of $K \subset S_k^1[x_0, x_{\mu+1}]$, the splines computed by the algorithm. Therefore the minimal approximation error is realized by the algorithm and a best approximation computed. \square

3. Realization, simplification and acceleration

As can be seen from Lemma 2.9 the computational cost and computing time grows exponentially with the number of data $\mu + 2$ causing a corresponding memory demand. It is advisable to restrain from computing and storing all (regular) position vectors since this would cause an exponential demand for memory. Instead, when a newly regular position vector is found, steps 3 and 4 should follow right away such that the position vector - once examined - can be deleted.

The algorithm is suitable for parallelization. Steps 3 and 4 can be executed for several regular position vectors in parallel.

Segments S_0 to S_r and their best approximations s_0, \dots, s_r could also be examined in parallel for intersection points. But this implementation is not advisable because it might lead to unnecessary computations: If the approximations s_0, s_1, \dots, s_r on segments S_0, S_1, \dots, S_r are computed in parallel and for example s_0 and s_1 do not intersect in (x_{i_0}, x_{i_0+1}) , then the position vector under examination can already be dismissed and the computation of s_2, \dots, s_r (for $r \geq 2$) in the parallel realization of the algorithm would be an additional and unnecessary effort.

In this respect more effective is sequential programming, starting step 4, computing s_0 and s_1 , and then check whether they intersect in (x_{i_0}, x_{i_0+1}) . In analogy a best approximation s_{l+2} on the next segment S_{l+2} is determined only if s_l and s_{l+1} have been found to intersect in (x_{i_l}, x_{i_l+1}) . Otherwise, the position vector currently under investigation is sorted out and computing further approximations related to this position vector can be skipped.

Furthermore, the algorithm should be programmed such that if the program terminates prematurely because of time limitations at least the best approximation found so far is saved and handed over to the main program. If position vectors of good approximations are dealt with early the consequences of a premature termination can be further limited.

If there is prior knowledge about the position of knots of a best approximation related position vectors should be examined with priority.

Which clues for the optimal position of knots can be deduced from the data? Besides the divided differences of first order

$$\Delta_j^{(1)} := \frac{f_{j+1} - f_j}{x_{j+1} - x_j}, \quad 0 \leq j \leq \mu$$

the (central) second order divided differences

$$\Delta_j^{(2)} := \frac{f_{j+1} - 2f_j + f_{j-1}}{(x_{j+1} - x_j)^2}, \quad 1 \leq j \leq \mu.$$

can be helpful here. The knots of a best-approximating first-order spline could be expected in the neighborhood of data abscissae x_j where the $\Delta_j^{(1)}$ vary heavily and the absolute values of the $\Delta_j^{(2)}$ are "large". Related positions vectors should therefore be examined with priority.

4. Numerical examples and applications

In this section we present numerical examples to illustrate the performance of Algorithm 2.10. For each example we list the knots and the approximation error and present the solutions in graphical form, but refrain from enumerating the coefficients of the best approximations in all examples.

Example 4.1. Let $\mu = 15$, $x_i := i$ for $i = 0, 1, \dots, 16$, $f_i := 1$ for $i = 0, 1, \dots, 7, 9, \dots, 16$ and for $f_8 := 2$ be given. For $k = 1, 2, 3, 4, 5$ Algorithm 2.10 computes best approximations $s_k \in S_k^1[t_0, t_{16}]$ with knots t_1, \dots, t_k given in the following table:

	t_1	t_2	t_3	t_4	t_5	$\ F - s_k(X)\ _2$
$k = 1$	8					0.87586
$k = 2$	7	8				0.78881
$k = 3$	7	8	9			0
$k = 4$	4	7	8	9		0
$k = 5$	1	2	7	8	9	0

For $k = 1, 2, 3$ s_k has exactly k proper knots. To approximate the data for $k \geq 4$ only 3 proper knots are needed and the remaining $k - 3$ knots are not proper knots; see Fig. 3.




Figure 3: Best approximation from Example 4.1.

Example 4.2. The so-called Titanium Heat Data have been discussed in literature and approximated from a number of approximation families; e.g. [5] and [6]. The data describe a property of titanium as function of temperature. The measuring points are equidistant in the interval [595, 1075]. The least-squares approximation with broken lines delivers a good approximation as can be seen in Fig. 4 and the following table::

k	$\ F - s_k(X)\ _2$	i	1	2	3	4	5
3	0.2632	t_i	858.4883	897.8327	940.2917		
		$s_k(t_i)$	0.7642	2.3065	0.6659		
4	0.1875	t_i	831.4392	866.8552	897.5429	940.2917	
		$s_k(t_i)$	0.7074	1.0396	2.3177	0.6659	
5	0.1349	t_i	831.4392	866.8552	898.3019	930.6129	958.3397
		$s_k(t_i)$	0.7074	1.0396	2.3494	0.9535	0.6153

Example 4.3. We look at the equidistant data abscissae $x_i = -1 + \frac{i}{10}$, $i = 0, 1, \dots, 20$ and the convex values $f_i := x_i^2$, $i = 0, 1, \dots, 20$. Approximation of the data (x_i, f_i) by a first-order spline with $k = 5$ free knots preserves the convexity of the data as can be seen in Fig. 5.




Figure 4: Best approximation of the Titanium Heat Data by first-degree splines with $k = 3$ bzw. $k = 5$ free knots respectively.

Example 4.4. For $k = 2, 3$ the data (x_i, f_i) from Fig. 6 were approximated by broken lines from $S_k^1[x_0, x_{71}]$. The values f_i are the arithmetic mean of measurements of forces measured in approximately equidistant time intervals when a sub-way switch ist set.

For $k = 2$ a best approximation s_2 was computed with the (interior) knots $t_1 = 0.77277$ and $t_2 = 1.83478$ and approximation error $\|F - s_2(X)\|_2 = 686.36122$. Raising the number of free knots to $k = 3$ lowers the approximation error to $\|F - s_3(X)\|_2 = 460.5584$ where the computed best approximation s_3 has the three (interior) knots $t_1 = 0.25851, t_2 = 0.87598$ and $t_3 = 1.81896$.

Example 4.5. Fig. 7 shows the best approximations from $S_7^1[x_0, x_{19}]$ and from $S_{10}^1[x_0, x_{19}]$ to $\mu + 2 = 20$ data points (x_i, f_i) with $x_i := i + 1, i = 0, 1, 2, \dots, 20$. In computing the best approximation in $S_7^1[x_0, x_{19}]$ Algorithm 2.10 inspected 6 724 520 position vectors with $1 \leq p(1) < p(2) < \dots < p(10) \leq 2\mu - 1$, only 795 455 of which are regular. For the approximation from $S_{10}^1[x_0, x_{19}]$ the algorithm inspected 183 579 396 position vectors with $1 \leq p(1) < p(2) < \dots < p(10) \leq 2\mu - 1$, of which 1 256 465 were regular.

Next, we present a medical application which originally motivated the development of Algorithm 2.10. The medical objective is to determine with an automated procedure the lowest concentration of an antibiotic necessary to kill (a defined high proportion of) the targeted bacteria, the so-called *minimal bactericidal concentration* (MBC). In clinical applications this helps to fight germs effectictely whith minimal unwanted side effects.

The measurements are determined as follows: An inital concentration κ_0 of the antibiotic is halved in each step. Germs are then exposed to each dilution $\kappa(j) := \kappa_0 \cdot 2^{-j}, j = 0, \dots, z, z \in \mathbb{N} \setminus \{0\}$. The effect of concentration $\kappa(j)$ on the viability of the germs ist measured by a flourescence method which allows to derive the values

$$\varphi_j := \frac{\text{number of surviving germs after exposition}}{\text{total number of germs before exposition}}$$

for $j = 0, \dots, z$.

This gives us $z + 1$ data points $(\kappa(0), \varphi_0), \dots, (\kappa(z), \varphi_z)$. For better representation instead of concentration $\kappa(j)$ the logarithmic values $\log(\kappa(j))$ are used, that ist, we make use of the relation $j = -\frac{\log(\kappa(j)) - \log(\kappa_0)}{\log(2)}$ and plot the pairs

$$(0, \varphi_0), (1, \varphi_1), \dots, (z, \varphi_z).$$




Figure 5: The best approximation to convex data is convex again in Example 4.3.




Figure 6: Best approximation from Example 4.4.

See Fig. 8 for a typical graph of the data and a best approximating first-degree spline $s^* \in S_2^1[x_0, x_z]$ with knots t_1 and t_2 . The knot t_1 is interpreted from the medical perspective as *minimal bactericidal concentration*

$$\text{MBC} := \kappa(t_1) = \kappa_0 \cdot 2^{-t_1}$$

Also of medical interest is knot t_2 from which

$$\text{MIC} := \kappa(t_2) = \kappa_0 \cdot 2^{-t_2},$$

is derived, the *minimal inhibitory concentration*, the lowest concentration with noticeable bactericidal effect; see [1] for more details.

In the following examples the initial concentration was $\kappa_0 := 256 = 2^8$, when $z = 19$ and $\kappa_0 := 128 = 2^7$, when $z = 18$. The following numerical example shows typical data and a typical best approximation $s^* \in S_2^1[x_0, x_z]$ with monotonically increasing s^* .

Example 4.6. With $x_i := i$ for $i = 0, 1, \dots, 19$ the following medical values f_i must be approximated:

i	0	1	2	3	4	5	6	7	8	9	10
f_i	3.6273	3.381	3.0339	2.8414	2.7507	2.9006	2.941	2.9986	3.2127	3.8381	8.2629




Figure 7: Best approximations $s_7 \in S_7^1[1, 20]$ and $s_{10} \in S_{10}^1[1, 20]$ to the 20 data points (x_i, f_i) from Example 4.5.

i	11	12	13	14	15	16	17	18	19
f_i	37.7363	84.0146	94.7914	98.7679	97.0424	98.0432	95.5602	99.0313	100

As best approximation from $S_2^1[x_0, x_{19}]$ Algorithm 2.10 computed the broken line

$$s^*(x) = \begin{cases} 0.2368x + 2.43313 & , x < 10.28981 \\ 46.2783x - 471.325 & , 10.28981 \leq x < 12.25123 \\ 0.52394x + 89.22211 & , x \geq 12.25123 \end{cases}$$

with the interior knots $t_1 = 10.2898$ and $t_2 = 12.2512$ and the approximation error $\|F - s^*(X)\|_2 = 5.7246$; see Fig. 8. This gives us the following values for MIC and MBC

$$\text{MBC} = 2^{8-t_1} = 0.2045 \quad \text{and} \quad \text{MIC} = 2^{8-t_2} = 0.0525.$$




Figure 8: Best approximation s^* from Example 4.6.

As seen in the above example the two knots t_1 and t_2 of a best approximation partition the medical data into three segments. In many cases the values are slightly monotonically increasing in the first and the last segment and ascending rapidly in the middle segment.

It may happen that no measurements are available from the middle segment. The following examples show that not always two (proper interior) knots are needed for a best approximation and the solutions need not to be unique.

Example 4.7. For $i = 0, 1, \dots, 18$ the medical data (x_i, f_i) are given by $x_i := i$ and the values f_i from the following table:

i	0	1	2	3	4	5	6	7	8	9
f_i	4.8245	5.0786	5.7781	6.105	5.9493	6.0516	5.589	5.5087	5.2563	4.5123
i	10	11	12	13	14	15	16	17	18	
f_i	97.8802	96.3044	95.6139	98.974	95.9425	96.0353	97.0482	98.5606	100	

As best approximation from $S_2^1[x_0, x_{18}]$ Algorithm 2.10 computed the first-degree spline

$$s^*(x) = \begin{cases} -0.026416x + 5.58421 & , x < 9 \\ 91.01492x - 813.78779 & , 9 \leq x < 10 \\ 0.25296x + 93.83177 & , x \geq 10 \end{cases}$$

with the data knots $t_1 = 9$ and $t_2 = 10$, see Fig. 9. The associated approximation error is $\|F - s^*(X)\|_2 = 4.24581$ and we compute $MBC = 2^{7-t_1} = 0.25$ and $MIC = 2^{7-t_2} = 0.125$.

Algorithm 2.10 computed just one best approximation, namely s^* , but there exist infinitely many best approximations to the given data: Each of the functions

$$s(y_1, y_2, x) := \begin{cases} s_1(x) & , x \leq y_1 \\ s_2(x) & , y_1 < x < y_2 \\ s_3(x) & , x \geq y_2 \end{cases} \quad := \begin{cases} -0.026416x + 5.58421 & , x \leq y_1 \\ \frac{s_3(y_2) - s_1(y_1)}{y_2 - y_1} (x - y_1) + s_1(y_1) & , y_1 < x < y_2 \\ 0.25296x + 93.83177 & , x \geq y_2 \end{cases}$$

is a best approximation from $S_2^1[x_0, x_{18}]$ to the data $(x_0, f_0), \dots, (x_{18}, f_{18})$ for arbitrary y_1, y_2 with $t_1 \leq y_1 < y_2 \leq t_2$. As limit case we get the *discontinuous* functions

$$s(y, x) := \begin{cases} -0.026416x + 5.58421 & , x \leq y \\ 0.25296x + 93.83177 & , x > y \end{cases}$$

with jump discontinuity in y for arbitrary $y \in (t_1, t_2)$ as best approximations to the given data.




Figure 9: Best approximations s^* and $s(9.4, 9.6, x)$ from Example 4.7.

Example 4.8. For $i = 0, 1, \dots, 18$ we look at the data (x_i, f_i) from Example 4.7 except that now $f_9 := 7.5123$ and $f_{10} := 97.8802$. As best approximation from $S_2^1[x_0, x_{18}]$ to the data pairs $(x_0, f_0), \dots, (x_{18}, f_{18})$

thus defined we get from Algorithm 2.10 the spline

$$s^*(x) = \begin{cases} 0.043098x + 5.39884 & , x < 8.98057 \\ 88.84909x - 792.12948 & , 8.98057 \leq x < 10 \\ 0.25296x + 93.83177 & , x \geq 10 \end{cases}$$

with the interior knot $t_1 = 8.98057$, the data knot $t_2 = 10$ and the minimal approximation error $\|F_2 - s_2^*(X)\|_2 = 4.11872$, see Fig. 10. Here again, arbitrarily many solutions $s(y_2, t) \in S_2^1[t_0, t_{18}]$ can be derived from s^* for all of which

$$s(y_2, x_9) = s^*(x_9) = f_9 = 7.5123$$

must hold. This implies that for the knots y_1 and y_2 the inequality $t_1 \leq y_1 < x_9 < y_2 \leq t_2$ must be valid. Because of $s(y_2, x_9) = f_9$ only one of the two knots can be freely chosen, the other one depends on that choice. If e.g. y_2 with $x_9 < y_2 < t_2$ ist chosen, then the splines

$$s(y_2, t) := \begin{cases} s_2(t) \\ s_2(t) \\ s_3(t) \end{cases} := \begin{cases} 0.043098x + 5.39884 & , x \leq y_1 \\ \frac{f_9 - s_3(y_2)}{x_9 - y_1}(x - x_9) + f_9 & , y_1 < x < y_2 \\ 0.25296x + 93.83177 & , x \geq y_2 \end{cases}$$

also are best approximations, where y_1 ist determined as solution of $s_1(y_2) = s_2(y_2)$.




Figure 10: The best approximations s^* , $s(9.05, x)$ and $s_2(9.5, x)$ from Example 4.8.

Measurement data not always meet the pattern of Example 4.6. Whether the interpretation of computed knots als MIC and MBC ist still a good idea must be decided from the medical perspective. In the next example we focus on such a situation.

Example 4.9. For $i = 0, 1, \dots, 19$ the medical data pairs (x_i, f_i) are listed in the following table:

i	0	1	2	3	4	5	6	7	8	9
f_i	3.0354	3.1654	3.0862	3.0564	2.9804	2.9632	2.8198	3.1239	3.0576	2.9828
i	10	11	12	13	14	15	16	17	18	19
f_i	3.1498	3.5877	4.0296	6.6481	9.829	12.1237	30.1584	70.2245	89.7225	100

As best approximation from $S_2^1[t_0, t_{19}]$ Algorithm 2.10 computed the spline

$$s^*(x) = \begin{cases} 0.40661t + 1.30283 & , t < 15.43646 \\ 40.0661t - 610.8992 & , 15.43646 \leq t < 17.30953 \\ 10.2775t - 95.2725 & , t \geq 17.30953 \end{cases}$$

with the two (interior) knots $t_1 = 15.43646$ and $t_2 = 17.30953$ and the approximation error $\|F - s^*(X)\|_2 = 7.69589$, see Fig. 11. This leads to $MBC = 2^{8-t_1} = 0.00577$ and $MIC = 2^{8-t_2} = 0.00158$.




Figure 11: The best approximation s^* from Example 4.9.

References

- [1] Bär, W., Både-Schumann, U., Krebs, A., Cromme, L. J.: Rapid method for detection of minimal bactericidal concentration of antibiotics. *Journal of Microbiological Methods*, 77, 85-89, (2009)
- [2] Cromme, L. J.: Manifolds with cusps. *Mathem. Nachrichten*, 142, 219-234 (1989)
- [3] Cromme, L. J.: A unified approach to differential characterizations of local best approximations for exponential sums und splines. *J. of Approximation Theory*, 36(4), 294-303 (1982)
- [4] Cromme, L. J., Kunath, J.: Discrete approximation by first-degree splines with free knots. Preprint at <https://arxiv.org> (2017).
- [5] de Boor, C.: *A practical guide to splines*, Springer, New York (1978)
- [6] Jupp, D. L. B.: Approximation to data by splines with free knots. *SIAM J. Numer. Anal.*, 15(2), 328-343 (1978)

Appendix A: Theorem 12 from [4]

Theorem 12. Let $\mu \geq k + 1 \geq 2$ and $1 \leq p \leq \infty$. For given data $(x_0, f_0), \dots, (x_{\mu+1}, f_{\mu+1})$ with $x_0 < x_1 < \dots < x_\mu < x_{\mu+1}$ and $f_0, \dots, f_{\mu+1} \in \mathbb{R}$ exists a best approximation $s^* \in S_k^1[x_0, x_{\mu+1}]$

$$\|f - s^*\|_{p,X} = \inf_{s \in S_k^1[x_0, x_{\mu+1}]} \|f - s\|_{p,X},$$

with the following additional features where t_1, \dots, t_k with $t_0 := x_0 < t_1 < \dots < t_k < t_{k+1} := x_{\mu+1}$ denote the knots of s^* :

(a) There are no knots in the boundary regions. More precisely:

$$x_0 < x_1 \leq t_1 < \dots < t_k \leq x_\mu < x_{\mu+1} \quad .$$

(b) Data abscissae neighboring to interior knots are not knots.

(c) Between two (not necessarily neighboring) interior knots of s^* lie at least two data abscissae which are not knots of s^* .

(d) On or between neighboring knots lie at least two data abscissae:

$$\forall j, 0 \leq j \leq r : \exists i, 0 \leq i \leq \mu : t_j \leq x_i < x_{i+1} \leq t_{j+1} \quad .$$

(e) If an interior knot is situated between two neighboring data abscissae, then no additional knot lies on or between these data abscissae. That is, the proposition " t_j is a knot of s^* with $x_q < t_j < x_{q+1}$ " implies $t_{j-1} < x_q$ and $t_{j+1} > x_{q+1}$.

(f) Let t_j be an interior knot. Then in each of the intervals $(-\infty, t_j)$ and (t_j, ∞) there is a data abscissa from $\{x_1, \dots, x_\mu\}$ which is not a knot.

(g) For $p < \infty$ we have: Between an interior and a neighboring data knot of s^* lies either exactly one data abscissa x_q which is then reproduced ($t_j < x_q < t_{j+1}$, $s^*(x_q) = f_q$) or there exist at least two data abscissae x_q, x_{q+1} between the knots ($t_j < x_q < x_{q+1} < t_{j+1}$).

(h) All interior knots are proper knots, that is, the first derivative is discontinuous in all interior knots.